



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2011

---

## **High-dose steroid treatment increases free water transport in peritoneal dialysis patients**

de Arteaga, J ; Ledesma, F ; Garay, G ; Chiurchiu, C ; de la Fuente, J ; Douthat, W ; Massari, P ;  
Terry, S ; Devuyst, O

**Abstract:** The water channel aquaporin-1 (AQP1) is the molecular counterpart of the ultrasmall pore that mediates free water transport during peritoneal dialysis (PD). Proof-of-principle studies performed in rats have shown that treatment with corticosteroids upregulates the expression of AQP1 in the peritoneal capillaries, causing a significant increase in free water transport. Whether such a beneficial effect could be observed in end-stage renal disease patients treated by PD remains unknown. Peritoneal transport parameters were evaluated in three patients on PD, shortly before and after living-donor renal transplantation and treatment with high-dose methylprednisolone (1.0-1.2 g/m<sup>2</sup>). As compared with pre-transplantation values, the post-transplantation test revealed an 2-fold increase in the sodium sieving and ultrasmall pore ultrafiltration volume, suggesting an effect on AQP1 water channels. In contrast, there was no change in the parameters of small solute transport. The direct involvement of AQP1 in these changes is suggested by the expression of glucocorticoid receptors in the human peritoneum and the presence of conserved glucocorticoid response elements in the promoter of the human AQP1 gene.

DOI: <https://doi.org/10.1093/ndt/gfr533>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-53203>

Journal Article

Accepted Version

Originally published at:

de Arteaga, J; Ledesma, F; Garay, G; Chiurchiu, C; de la Fuente, J; Douthat, W; Massari, P; Terry, S; Devuyst, O (2011). High-dose steroid treatment increases free water transport in peritoneal dialysis patients. *Nephrology, Dialysis, Transplantation*, 26(12):4142-4145.

DOI: <https://doi.org/10.1093/ndt/gfr533>

NDT-01191-2011R1

*Exceptional Case*

**HIGH-DOSE STEROID TREATMENT INCREASES FREE-WATER TRANSPORT  
IN PERITONEAL DIALYSIS PATIENTS**

Javier de Arteaga<sup>1,2\*</sup>, Fabian Ledesma<sup>1,2</sup>, Gabriela Garay<sup>1,2</sup>, Carlos Chiurchiu<sup>1,2</sup>, Jorge de la  
Fuente<sup>1,2</sup>, Walter Douthat<sup>1,2</sup>, Pablo Massari<sup>1,2</sup>, Sara Terryn<sup>3</sup>, and Olivier Devuyst<sup>3,4\*</sup>

<sup>1</sup> Servicio de Nefrología, Hospital Privado de Córdoba, Universidad Católica de Córdoba;

<sup>2</sup> Fundación Nefrológica de Córdoba, Córdoba, Argentina;

<sup>3</sup> Division of Nephrology, Université catholique de Louvain Medical School, Brussels,  
Belgium;

<sup>4</sup> Institute of Physiology, Zurich Center for Integrative Human Physiology, Klinik für  
Nephrologie, University of Zurich, Zurich, Switzerland

**\*Correspondence:** Prof. Dr. Olivier Devuyst  
Email: [olivier.devuyst@uzh.ch](mailto:olivier.devuyst@uzh.ch)  
or  
Prof. Dr. Javier de Arteaga  
Email: [javierdearteaga@gmail.com](mailto:javierdearteaga@gmail.com)

Word count: 1000

Short title: Free-water transport in peritoneal dialysis

## ABSTRACT

The water channel aquaporin-1 (AQP1) is the molecular counterpart of the ultrasmall pore that mediates free-water transport during peritoneal dialysis. Proof-of-principle studies performed in rats have shown that treatment with corticosteroids upregulates the expression of AQP1 in the peritoneal capillaries, causing a significant increase in free-water transport. Whether such a beneficial effect could be observed in end-stage renal disease patients treated by peritoneal dialysis remains unknown. Peritoneal transport parameters were evaluated in three patients on peritoneal dialysis, shortly before and after living-donor renal transplantation and treatment with high-dose methylprednisolone (1.0 to 1.2 gr/m<sup>2</sup>). As compared with pre-transplantation values, the post-transplantation test revealed a ~2-fold increase in the sodium sieving and ultrasmall pore ultrafiltration volume, suggesting an effect on AQP1 water channels. In contrast, there was no change in the parameters of small solute transport. The direct involvement of AQP1 in these changes is suggested by the expression of glucocorticoid receptors in the human peritoneum and the presence of conserved glucocorticoid response elements in the promoter of the human *AQP1* gene.

**Keywords:** aquaporin-1, endothelium, peritoneal capillaries, sodium sieving

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The capacity for ultrafiltration (UF) across the peritoneal membrane is a major predictor of outcome and mortality for end-stage renal disease patients treated with peritoneal dialysis (PD) (1,2). The major barrier for the transport of water and solutes across the peritoneal membrane is the endothelium lining peritoneal capillaries. Modeling of peritoneal transport has predicted that endothelial cells contain ultrasmall pores (radius < 3 Å) that facilitate the osmotic transport of water during a hypertonic dwell (3). The ultrasmall pores account for half of the UF and their patency explains the “sodium sieving”, i.e. the marked fall of the dialysate to plasma ratio of sodium during the first hour of PD (3). Several lines of evidence have demonstrated that the water channel aquaporin-1 (AQP1), which is abundantly expressed in the endothelial cells lining peritoneal capillaries and veinules (4), is the molecular counterpart of the ultrasmall pores (5, for review). Studies in *Aqp1*-null mice demonstrated that AQP1 facilitates water transport and mediates approximately half of the UF during crystalloid osmosis (6,7). Accordingly, increasing the expression of AQP1 in the peritoneum could improve water transport in PD patients. Based on the presence of glucocorticoid response elements (GREs) in the promoter of the murine AQP1 gene (8), Stoenoiu et al. demonstrated that treatment with corticosteroids increases the expression of AQP1 in the peritoneal capillaries, mirrored by increased water transport and UF, without affecting the transport of small solutes (9). Thus far, it is unknown whether a similar effect could be evidenced in patients undergoing PD. We report the serial determination of water and solute transport parameters across the peritoneal membrane in three PD patients that received high-dose glucocorticoids as standard care after a living-donor renal transplantation.

## PATIENTS AND METHODS

The clinical features of the three PD patients are detailed in Table 1. *Patient 1* is a 11 year-old girl who presented hemolytic uremic syndrome at 18 months of age, progressing to ESRD at age 10. She was initially treated by CAPD for 6 months, followed by APD for 4 months. *Patient 2* is a 54 year-old male with autosomal dominant polycystic kidney disease (ADPKD) progressing to ESRD at age 52. He was treated by CAPD for 9 months before transplantation. *Patient 3* is a 6 year-old boy who reached ESRD at age 5 for congenital vesico-ureteral reflux. He was treated by CAPD for 5 months. The course of PD was uncomplicated in the three patients. They underwent a living-donor renal transplantation after a mean of 8 months of PD (range: 5 to 10 months). The renal transplantation was uneventful, with immediate graft function in the three cases. The detail of cumulative methylprednisolone dose and immunosuppressive treatment is given in Table 1. The peritoneal function was assessed in the three patients using the Mini-Peritoneal Equilibration Testing (Mini-PET) (10) performed before (range: 1 day to 3 months) and after (range: 7 to 39 days) renal transplantation. The sodium concentration was measured by indirect ion selective analysis (Modular Analytics Evo 900, calibrated by ISE compensator; Roche Diagnostics, Buenos Aires, Argentina). The study was approved by the Ethical Review Board and informed consent was obtained.

The expression of the glucocorticoid receptor (GR) mRNA was assessed in human peritoneal biopsies using reverse transcription polymerase chain reaction (RT-PCR) (9). Total RNA was extracted from two human peritoneal biopsies (obtained a time of catheter removal in PD patients, Ref. 4) using Aurum<sup>TM</sup> Total RNA Tissue Kit (Bio-Rad, Hercules, CA), following the manufacturer's protocol, followed by DNase I treatment to eliminate genomic DNA contamination. Total RNA was extracted from human erythroleukemia (HEL) cells (positive control for GR expression) using RNAqueous<sup>R</sup>-Micro kit (Ambion). One microgram

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

of RNA was used to perform the reverse transcriptase reaction with iScript™ cDNA Synthesis Kit (Bio-Rad). RT-PCR detection was performed with FastStart Taq DNA Polymerase (Roche) for 32 cycles, and PCR products visualized on a 1.5% agarose gel with GelRed dye (Sigma). The primers were designed using Beacon Design 2.0 (Premier Biosoft International, Palo Alto, CA) : GAPDH (forward 5'-3' : GGG GCT CTC CAG AAC ATC AT; and reverse 5'-3': TCT AGA CGG CAG GTC AGG T) and GR (forward 5'-3': ACA CTG CCC CAA GTG AAA AC; and reverse 5'-3': CCA TGA ACA GAA ATG GCA GA).

Immunostaining was performed on paraffin-embedded sections of human peritoneal biopsies by using a sequential staining protocol (4). Sections were incubated with 3% serum for 20 min before adding anti-von Willebrand factor ( $\alpha$ -vWF, DAKO, Denmark) diluted in phosphate-buffered saline (PBS) containing 2% BSA for 1h. After washing, sections were incubated with Alexa633-labelled secondary anti-IgG antibodies (Invitrogen, Belgium) for 45 min. The secondary staining was performed by adding anti-glucocorticoid receptor ( $\alpha$ -GR, Abcam, Belgium) and labelled with Alexa488-labelled secondary anti-IgG antibodies before mounting in Prolong Gold Anti-fade reagent (Invitrogen, Belgium). Sections were viewed under a Zeiss LSM510Meta Confocal microscope (Zeiss, Belgium), using x20 Plan-Apochromat or x63/1.4 Plan-Apochromat oil-immersion objectives (Zeiss).

The 3kb promoter sequence of the human *AQP1* gene was analyzed for putative GRE using Transcription Element Search Software (TESS; <http://www.cbil.upenn.edu/cgi-bin/tess/tess>).

## RESULTS AND DISCUSSION

The evolution of the peritoneal transport parameters as assessed by the Mini-PET is shown on Table 1. As compared with pre-transplantation, the post-transplantation PET evidenced an approximately 2-fold increase in the sodium sieving (and its surrogate marker the absolute dip of dialysate sodium concentration) (11), paralleled by a similar increase in the ultrasmall pore-specific UF. In contrast, the small solute transport rate, as assessed by the D/P creatinine and small pore-specific UF was unchanged (Fig. 1A, Table 1). By analogy with results obtained in the rat (9), the potential role of AQP1 was substantiated by expression studies demonstrating the expression of the GR in the capillary endothelium of the human peritoneum, and the *in silico* identification of conserved GREs in the promoter of the *AQP1* gene (Fig. 1B-D).

These data are the first to document a major increase in the parameters of free-water transport in PD patients following high-dose glucocorticoid administration in the context of renal transplantation. To the best of our knowledge, there are no previous reports on the effect of renal transplantation *per se* on PD transport parameters. Several points argue in favour of a specific effect of the glucocorticoids on AQP1. First, the time-course of the effect, similarly observed in the three patients who had all a short course of uncomplicated PD followed by a living-donor renal transplantation with immediate renal function. Second, the fact that the changes are restricted to the very parameters – sodium sieving, sodium dip and ultrasmall pore-specific UF – that reflect the AQP1-mediated water transport, whereas there is no effect on the D/P creatinine parameter assessing small solute transport. The lack of effect of small solutes is important when considering that exposure to mycophenolate mofetil (MMF) can, at least *in vitro*, inhibit angiogenesis (12,13) and thus decrease the exchange area. Third, the fact that no other external factor other than high-dose glucocorticoids has been reported to specifically affect free-water transport and expression of AQP1. Fourth, we provide evidence

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

for multiple GREs in the promoter of the *AQP1* gene and for a strong expression of the GR in the endothelium lining peritoneal capillaries, i.e. in the very cells where AQP1 mediates osmotic water transport during PD. We fully acknowledge the limitations of this preliminary report, including the small number of patients, with only one adult, not very representative of the PD population, and the lack of low-dose steroid controls assessing the effect of transplantation (e.g. removing uremia, inducing acute changes in body parameters, and putting the peritoneal membrane at rest). These limitations are balanced by the specificity of the changes in free-water transport and the lack of effect on other transport parameters (Figure 1). We should also keep in mind the possibility of a role for the interstitium in the water-only fluid pathway (14), although the time-course for a dramatic reorganization of the interstitium (except for a reduction in the degree of oedema) seems to be too short for such an explanation of the observed changes.

In conclusion, these data provide the first line of evidence supporting the effect of corticosteroids on specific parameters of AQP1-mediated water transport in PD patients. Although these results have to be confirmed on a larger patient material, the specificity of the changes in free-water transport suggests that pharmacologic induction of AQP1 may help to increase water transport in patients treated by PD.



## ACKNOWLEDGEMENTS

We thank Y. Cnops, H. Debaix and J. Morelle for excellent technical help. The support of the Fonds de la Recherche Scientifique Médicale, the National Centre of Competence in Research (NCCR) Kidney.CH and the Fundacion Nefrológica de Córdoba is gratefully acknowledged.

## REFERENCES

1. Brown EA, Davies SJ, Rutherford P, et al. Survival of functionally anuric patients on automated peritoneal dialysis: the European APD Outcome Study. *J Am Soc Nephrol* 2003; 14: 2948-2957
2. Brimble KS, Walker M, Margetts PJ, Kundhal KK, Rabbat CG. Meta-analysis: peritoneal membrane transport, mortality, and technique failure in peritoneal dialysis. *J Am Soc Nephrol* 2006; 17: 2591-2598
3. Rippe B, Stelin G, Haraldsson B. Computer simulations of peritoneal fluid transport in CAPD. *Kidney Int* 1991; 40: 315-325
4. Devuyst O, Nielsen S, Cosyns JP, Smith BL, Agre P, Goffin E. Aquaporin-1 and endothelial nitric oxide synthase expression in capillary endothelia of human peritoneum. *Am J Physiol* 1998; 275: H234-H242
5. Nishino T, Devuyst O. Clinical application of aquaporin research: Aquaporin-1 in the peritoneal membrane. *Pflügers Arch* 2008; 456: 721-727
6. Yang B, Folkesson HG, Yang J, Matthay MA, Ma T, Verkman AS. Reduced osmotic water permeability of the peritoneal barrier in aquaporin-1 knockout mice. *Am J Physiol* 1999; 276: C76-C81
7. Ni J, Verbavatz JM, Rippe A, et al. Aquaporin-1 plays an essential role in water permeability and ultrafiltration during peritoneal dialysis. *Kidney Int* 2006; 69: 1518-1525
8. King LS, Nielsen S, Agre P. Aquaporin-1 water channel protein in lung: Ontogeny, steroid-induced expression, and distribution in rat. *J Clin Invest* 1996; 97: 2183-2191
9. Stoenoiu MS, Ni J, Verkaeren C, et al. Corticosteroids induce expression of aquaporin-1 and increase transcellular water transport in rat peritoneum. *J Am Soc Nephrol* 2003; 14: 555-565
10. La Milia V, Limardo M, Virga G, Crepaldi M, Locatelli F. Simultaneous measurement of peritoneal glucose and free water osmotic conductances. *Kidney Int* 2007; 72: 643-650
11. La Milia V, Pozzoni P, Virga G, et al. Peritoneal transport assessment by peritoneal equilibration test with 3.86% glucose: A long-term prospective evaluation. *Kidney Int* 2006; 69: 927-933
12. Duncan FJ, Wulff BC, Tober KL, et al. Clinically relevant immunosuppressants influence UVB-induced tumor size through effects on inflammation and angiogenesis. *Am J Transplant* 2007; 7:2693-2703
13. Koehl GE, Wagner F, Stoeltzing O, et al. Mycophenolate mofetil inhibits tumor growth and angiogenesis in vitro but has variable antitumor effect in vivo, possibly related to bioavailability. *Transplantation* 2007; 83: 607-614

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

14. Rippe B, Venturoli D. Simulations of osmotic ultrafiltration failure in CAPD using a serial three-pore membrane/fiber matrix model. *Am J Physiol Renal Physiol* 2007; 292: F1035-F1043

For Peer Review

**FIGURE LEGEND**

*Figure 1. Transport parameters before and after renal transplantation and expression of the glucocorticoid receptor (GR) in the human peritoneum.*

**A.** Comparison of the transport parameters before and after renal transplantation.

A mini-PET was performed before and after transplantation in the three patients, who received methylprednisolone (1 g/m<sup>2</sup>) at time of transplantation. As compared with baseline values, the three patients show a ~ 2-fold increase in the sodium sieving and in the ultrasmall-pore specific (USP) ultrafiltration (mL) post-transplantation, whereas the transport of small solutes (D/P creatinine) remains unchanged. Individual values are given in Table 1.

**B.** Reverse transcription polymerase chain reaction (RT-PCR) demonstrating the expression of the GR (153 bp) in two peritoneal biopsies from patients on peritoneal dialysis and in the control human erythroleukemia (HEL) cells.

**C.** Immunolocalization of the GR in human peritoneum. Double immunostaining with anti-glucocorticoid receptor ( $\alpha$ -GR) and anti-von Willebrand factor ( $\alpha$ -vWF) antibodies show that the GR is expressed in endothelial cells lining capillaries and small vessels in the peritoneal membrane. Low magnification: bar: 50 $\mu$ m; detail: 10 $\mu$ m. m, mesothelium.

**D.** Organization of human aquaporin-1 (*AQP1*) gene promoter. The 3-kilobase (kb) proximal promoter of *AQP1* is schematically represented, including the transcription start site and the 16 predicted glucocorticoid response elements (GRE, vertical symbols).

**Table 1 R1:** Clinical and peritoneal equilibration test (Mini-PET) parameters in the three patients.

	Patient 1	Patient 2	Patient 3
Gender, age (years)	Female, 11	Male, 54	Male, 6
Cause of ESRD	Hemolytic uremic syndrome	ADPKD	Congenital reflux
Age at ESRD (years)	10	52	5
Duration of PD (months), modality	10, CAPD (6) then APD (4)	9, CAPD	5, CAPD
Residual diuresis on PD (mL per day)	500	1000	500
Time of pre-transplant mini-PET (days)	1 day	90 days	2 days
Methylprednisolone dose (gr per m <sup>2</sup> ) <sup>a</sup>	1.0	1.0	1.2
Other (immunosuppressive) drugs introduced per-transplantation	Daclizumab (1mg/kg) Tacrolimus) (0.15 mg/kg/day) MMF 1 gr/day	Tacrolimus: 0.15 mg/kg/day MMF: 2 gr/day	Daclizumab (1mg/kg) Tacrolimus: (0.15 mg/kg/day) MMF: 1 gr/day
Time to post-transplant Mini-PET (days)	7	20	39
Serum creatinine at posttransplant Mini-PET (mg/dL)	0.44	1.2	0.43

**Table 1 R1 (cont.)**

	Patient 1		Patient 2		Patient 3	
	Pre-TP	Post-TP	Pre-TP	Post-TP	Pre-TP	Post-TP
Sodium sieving (%) <sup>b</sup>	8.3	11.4	5.4	13.1	4.4	12.7
Sodium dip (mmol/L) <sup>c</sup>	11	15	7	17	6	17
Ultrafiltration USP (mL)	126	150	170	332	30	90
Ultrafiltration 1h (mL)	240	180	620	380	60	180
D/P creatinine	0.40	0.47	0.54	0.46	0.66	0.66
Volume infused (L/m <sup>2</sup> )	1.1	1.1	1.2	1.2	0.9	0.9

ADPKD, autosomal dominant polycystic kidney disease; APD, automated peritoneal dialysis; CAPD, continuous ambulatory peritoneal dialysis; ESRD, end-stage renal disease; MMF, mycophenolate mofetil; PD, peritoneal dialysis; PET, peritoneal equilibration test.

<sup>a</sup>, cumulative dose until post-transplantation Mini-PET; <sup>b</sup>, sodium sieving:  $(D_0/P_0 - D_{60}/P_{60})/D_0/P_0 \times 100$  (in %) (Ref. 7); <sup>c</sup>, sodium dip: absolute dip of dialysate sodium concentration at 60 min of the PET (mmol/L) (11).

D/P, dialysate-over-plasma ratio; Post-TP, post-transplantation; Pre-TP, pre-transplantation; USP, ultrafiltration

Figure 1.R1

A




Figure 1.R1

